



EP18EO – Culture of Safety

EP18EO: Provide eight of the most recent consecutive quarters of unit-or -clinic level nurse-sensitive, clinical indicator data to demonstrate outperformance of the mean, median or other measure of central tendency (benchmark provided by the vendor's national database). Data must be the most recent eight complete quarters available from the vendor (s). Select and report data for all applicable inpatient care units.

The required patient indicators for all inpatient care organizations include falls with injury, hospital-acquired pressure injuries (HAPI) stage 2 and above, and two others from the list:

- *Falls with Injury*
- *Hospital-acquired pressure injuries (HAPI) stage 2 and above*

Select two others:

- *Central-line associated bloodstream infections (CLABSI)*
- *Catheter-associated urinary tract infections (CAUTI)*
- *Clostridium difficile (CDIFF)*
- *Methicillin-resistant Staphylococcus aureus (MRSA)*
- *Venous thromboembolism (VTE)*
- *Peripheral Intravenous infiltration (PIV)*
- *Physical and sexual assault*
- *Device-related hospital-acquired pressure injuries (HAPI)*

Selected Nurse Sensitive Clinical Indicator Data

EP18EOa	Falls with Injury
EP18EOb	Hospital acquired pressure injuries (HAPI) stage 2 and above
EP18EOc	Central line associated bloodstream infections (CLABSI)
EP18EOd	Catheter associated urinary tract infections (CAUTI)